

**SUNSHINE SECONDARY SCHOOL**

**233/3  
CHEMISTRY  
PAPER 3  
PRACTICAL  
PRE MOCK 1 2017  
MARCH 2017  
2<sup>1</sup>/<sub>4</sub>HRS**

**NAME.....CLASS..... ADM NO.....**

**SIGNATURE.....INDEX..... DATE .....**

**INSTRUCTIONS**

- ❖ Answer all the questions on the spaces provided
- ❖ All working must be clearly shown where necessary
- ❖ Calculations and mathematical tables may be used

**FOR EXAMINERS USE ONLY**

<b>QUESTION</b>	<b>MAXIMUM SCORE</b>	<b>CANDIDATES SCORE</b>
1	11	
2	14	
3	15	
<b>TOTAL</b>	<b>40</b>	

1 You are provided with:

Solution M 0.2M hydrochloric acid,

Solution F containing 15.3g per litre of basic compound  $G_2X.H_2O$ .

You are required to determine the relative atomic mass of G.

**PRECEDURE:**

Place solution M in a burette ,pipette  $25\text{cm}^3$  of solution F into a  $250\text{cm}^3$  conical flask. Add two drops of methyl orange indicator and titrate. Record your results in the table below. Repeat the procedure two more times and complete table I.

Table I

	I	II	III
Final burette reading			
Initial burette reading			
Volume of solution M used ( $\text{cm}^3$ )			

(4mks)

a) What is the average volume of solution M.?

(1mk)

.....  
.....

b) Given that one mole of F reacts with 2moles of M. Calculate the;

i) Number of moles the basic compound,  $G_2X, 10H_2O$  in the volume of solution F used.

(2mks)

.....  
.....  
.....  
.....

ii) Concentration of solution F in moles per litre.

(2mks)

.....  
.....  
.....

iii) Relative formula mass of the basic compound,  $G_2X.10H_2O$ .

(1mk)

- .....  
.....  
.....
- iv) Relative atomic mass of G (Relative formula Mass of X=60 , atomic mass of H=1.0 , O=16.0). (1mk)
- .....  
.....  
.....

2 You are provided with:

- 1 1.89g of solid P, solid P is adiabatic acid  $H_2X$ .
- 2 0.5M Solution of the dibasic acid ,  $H_2X$  , Solution V.
- 3 Sodium hydroxide, Solution K.

You are required to determine:

- a)
  - i) the molar heat of solid P.
  - ii) the heat of reaction of one mole of the dibasic acid with sodium hydroxide.
- b) Calculate the heat of reaction of solid  $H_2X$  with aqueous sodium hydroxide.

### **PROCEDURE I.**

Place  $30\text{cm}^3$  of distilled water into a 100ml beaker. Measure the initial temperature of the water and record it in the table II below. Add all the solid P at once; stir the mixture carefully with the thermometer until all the solid dissolves. Measure the final temperature reached and records it in the table II

**Table II**

Final temperature ( $^{\circ}\text{C}$ )	
Initial temperature ( $^{\circ}\text{C}$ )	

(2mks)

- a) Determine the change in temperature  $\Delta T_1$  ( 1 mk)
- .....

b) Calculate the:

- i) Heat change when  $H_2X$  dissolves in water, (Assuming the heat capacity of the solution is  $4.2\text{Jg}^{-1}\text{K}^{-1}$  and density is  $1\text{g/cm}^3$ ) (2mks)
- .....  
.....  
.....

ii) Number of moles of the acid that were used. (Relative formula mass of  $H_2X$  is 126) (1mk)

.....  
.....  
.....

iii) Molar heat of solution  $\Delta H_1$  solution of the acid  $H_2X$ . (1mk)

.....  
.....  
.....

**PROCEDURE II.**

Place  $30\text{cm}^3$  of solution V into a  $100\text{cm}^3$  beaker. Measure the initial temperature and record it in table III below. Measure  $30\text{cm}^3$  of sodium hydroxide, solution K. Add all of the  $30\text{cm}^3$  of t of solution K at once to V in the beaker. Stir the mixture with the thermometer. Measure the final temperature reached and record it in table III.

**Table III.**

Final temperature ( $^{\circ}\text{C}$ )	
Initial temperature ( $^{\circ}\text{C}$ )	

( 1 ½ mks)

a) Determine the change in temperature,  $\Delta T_2$ . (½ mk)

.....

b) Determine the:

i) Heat change for the reaction (Assume the heat capacity of the solution is  $4.2\text{Jg}^{-1}\text{K}^{-1}$  and density is  $1\text{g/cm}^3$ ) (2mks)

.....  
.....  
.....  
.....


ii) Number of moles of the acid used ( $H_2X$ ). (1mk)

.....  
.....  
.....

iii) Heat of reaction,  $\Delta H_2$  of one mole of the acid  $H_2X$  with sodium hydroxide (1mk)

.....  
.....  
.....  
.....

d) Given that,



Calculate  $\Delta H_3$  for the reaction  $H_2X_{(s)} + 2OH^-_{(aq)} \longrightarrow 2H_2O_{(l)} + X^{2-}_{(aq)}$  (2mks)

.....  
.....  
.....  
.....  
.....

**QUESTION 3A**

a) You are provided with solid Q. Carry out the test indicated below and record your observations and deductions in the table below.

i) Place a spatula full of Q in a boiling tube. Add about 10cm<sup>3</sup> of distilled water and shake. Divide the resultant mixture into 4 portions.

Observation	Deductions
(1mk)	(1mk)

b) To the first portion add Barium nitrate solution followed by dilute nitric acid.

Observation	Deduction
(2mks)	(1mk)

c) To the second portion add 2-3 drops of sodium hydroxide till in excess.

Observation	Deduction
(2mks)	(1mk)

d) To the third portion add 2-3 drops of ammonia solutions till in excess.

Observation	Deduction
(2mks)	(1mk)

e) To the 4<sup>th</sup> portion add Pb (NO<sub>3</sub>)<sub>2</sub> solution

Observation	Deduction
(1mk)	(1mk)

### QUESTION 3B

You are provided with liquid X. You are required to carry the test below.

a) Place about 1cm<sup>3</sup> of substance X in a test tube. Add a small piece of sodium carbonate solid.

Observation	Deduction
(1mk)	(1mk)

b) To about 3cm<sup>3</sup> of X in a boiling tube, add acidified potassium chromate (vi) and warm.

Observation	Deduction
(1mk)	(1mk)

c) To about 3cm<sup>3</sup> of X add acidified potassium manganate (vii)

Observation	Deduction
(1mk)	(1mk)

**233/3**  
**CHEMISTRY**  
**PRACTICAL**  
**PAPER 3**  
**MARCH 2017**

**CONFIDENTIAL**

**PRE MOCK**

*Kenya Certificate of Secondary Education*

1. About 50cm<sup>3</sup> of solution V
2. About 50cm<sup>3</sup> of solution K
3. 1.89g of solid P oxalic acid accurately weighed and placed in a stopped container.
4. Thermometer
5. 5 dry test tubes in a test tube rack
6. Spatula
7. Bunsen burner
8. About 120cm<sup>3</sup> of solution M
9. About 90cm<sup>3</sup> of solution F
10. Liquid X-ethanol
11. Solid Q – 1g of solid zinc sulphate
12. Blue and red litmus papers.
13. A boiling tube.
14. Glass rod

**Access to:**

- a) Bunsen burner
- b) 2M sodium hydroxide with a dropper
- c) 2M Ammonium hydroxide
- d) Barium nitrate solution
- e) Lead nitrate solution
- f) Dilute nitric v acid
- g) Methyl orange with a dropper.
- h) Phenolphthalein indicator in a bottle dropper
- i) About 15cm<sup>3</sup> of liquid X
- j) Acidified potassium dichromate (VI) with a dropper.
- k) Acidified potassium manganate (vii)

1. Solution V is prepared by dissolving 63g of oxalic acid to make one litre of solution.
2. Solution K is prepared by dissolving 16g of sodium hydroxide pellets to make one litre of solution.
3. Solution M is prepared by dissolving 17cm<sup>3</sup> of concentrated hydrochloric acid to make one litre of solution.
4. Solution F is prepared by dissolving 15.3g of hydrated sodium hydrogen carbonate to make one litre of solution.