

Kenya Certificate of Secondary Education.

233/3 CHEMISTRY PAPER 3 PRACTICAL TIME: 2<sup>1</sup>/<sub>4</sub> HOURS.

#### **INSTRUCTIONS TO CANDIDATES.**

- Write your name and index number in the spaces provided above.
- Sign and write the date of exam in the spaces above.
- Answer **ALL** the questions in the spaces provided.
- You are not allowed to start working with the apparatus for the first 15 minutes of the 2<sup>1</sup>/<sub>4</sub> hours allowed time for the paper.
- Use the 15 minutes to read through the question paper and note the chemicals you require
- Mathematical tables and electronic calculators may be used.
- All working **MUST** be clearly shown where necessary.
- This paper consists of 6 printed pages. Candidates should check to ensure that all pages are printed as indicated and no questions are missing

#### FOR EXAMINER'S USE ONLY.

Question	Maximum score	Candidate's score
1	18	
2	12	
3	10	
Total score	40	

© 2014, Makindu district inter - secondary schools examination

233/3 Chemistry Practical

## **Question 1**

You are provided with:

- Dilute hydrochloric acid solution A •
- 0.1m sodium hydroxide solution B
- Kcaepaatpapeta.com 10g of a mixture of sodium hydrogen carbonate and sodium chloride per litre, solution C

You are required top determine;

- (i) Molarity of solution A in the second
- (ii) Percentage purity by mass of sodium hydrogen carbonate

# PROCEDURE 1

Fill the burette with solution A. Pipette 25cm3 of 0.1M sodium hydroxide solution B into a clean conical flask and and 2 drops of methyl orange indicator and titrate with solution A until a permanent pink colour occurs. Fill in the table below. Repeat the titration two more times and complete the table below.

# TABLE I

L.C.	¥.	1	2	3
or No.	Final burette reading (cm <sup>3</sup> )			
\$ <sup>0</sup>	Initial burette reading (cm <sup>3</sup> )			
	Volume of solution A used (cm <sup>3</sup> )			
(a)	Calculate the average volume of solution A used.			(4 Marks) (1 Mark)
(b)	Calculate the number of moles of hydrochloric acid so hydroxide solution B.	olution A tha	t reacted with 25c	m3 of sodium (2 Marks)
(c)	Calculate the concentration of solution A in moles per	tlitre		(1 Mark)
			•••••••••••••••••••••••••••••••••••••••	
		•••••		

## **PROCEDURE II**

Pipette 25cm3 of solution C into a conical flask, Titrate with solution A using 3 drops of methyl orange indicator. Record your results in table II below. **TABLE II** 

	1	2	3
Final burette reading (cm <sup>3</sup> )			
Initial burette reading (cm <sup>3</sup> )			
Volume of solution A used (cm <sup>3</sup> )			
			(

e. con	
() Charles the first of the second seco	Chemistry Paper 3 Practical
(a) Calculate the average volume of solution A used.	(1 Mark)
e <sup>20</sup>	••••••
	••••••
·····	
(b) Write an ionic equation for the reaction taking place between solution A and mixt	ure C. (1 Mark)
1 <sup>3</sup>	
(c) Calculate: $\sqrt{2^{2^{-1}}}$	
(i) Molarity of sodium hydrogen carbonate in moles per litre	(2 Marks)
······	
e <sup>e</sup>	
$\sqrt{e}$ (ii) Mass of sodium hydrogen carbonate in moles per litre	(1 Mark)
(ii) muss of sociality hydrogen encounce in mores per nuc	(1 mark)
\$ <sup>0</sup>	
	•••••
(iii) Mass of sodium chloride in the mixture	(1 Mark)

- (I) You are provided with solid F. Carry out the following tests and write down all the observations and Inferences.
  - (a) Place half spatula end full of solid F in a dry test tube. Heat gently then strongly until there is no further change. Test gas using a glowing splint.

Observations	Inferences
(1 mark)	(1 mark)

(b) Place the remaining solid F in a test tube, add about 10cm<sup>3</sup> of distilled water and shake vigorously. Divide the mixture into three portions.

(i) To the first portion, add 2M sodium hydroxide solution drop wise until in excess.

~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~	Observations	Inferences
	Let.	
	Stor.	
	x <sup>2</sup> e <sup>e</sup>	
	and .	
	×.	
	a a a a a a a a a a a a a a a a a a a	
	A* 6	
	er s	
	-2 <sup>0,1</sup>	
	(1 mark)	(1 mark)
(ii)	To the 2 <sup>nd</sup> portion, add ammonia solution dropwise	e till in excess.
	10 <sup>51</sup>	
Г	Observations	Inferences
	\$ <sup>5</sup>	
1 de		
L'ILO		
\$°		
	(1 mark)	(1 mark)
	(iii) To the $3^{ra}$ portion, add 4 drops of solution chlo	bride
Г	Observations	Inferences
	(1 mark)	(1 mark)

#### II

You are provided with liquid K, carry out the following tests on it.

(a) Place about one spatula end full of liquid K on a metallic spatula and ignite it in a Bunsen burner flame.

Observations	Inferences
(1 mark)	(1 mark)

			com	
		é	ç <b>e</b>	Chemistry Paper 3 Practical
(b)	To 2cm3 of liquid K add 3 drops of acid	ified KMn	$O_4$ solution.	× 1
	Observations		Inferences	
	to Visit www.freekc	<sup>gev</sup>		
	o <sup>ape</sup> r			
		½ mark)	( ½ mark)	
(c)	To 2cm3 of liquid K, add 3 drops of acid	dified K <sub>2</sub> C	$r_2O_7.$	
	Observations		Inferences	
	\$TEC Y			
NOT				
<sup>°</sup> O <sup>°</sup>				
	(	<sup>1</sup> /2 mark)	( <sup>1</sup> /2 mark)	

- 3. You are provided with solid E. Carry out tests below. Record your observations and inferences in the spaces provided.
  - (a) Put about one half of solid E in a dry test tube and heat it strongly. Test for any gas produced using litmus paper.

Observations	Inferences
( 2 mark)	( 2 mark)

(b) Dissolve the rest of the solid E in 10cm3 of distilled water in boiling tube. Divide solution into 3 portions.

(i) To a first portion in test-tube, add aqueous sodium hydroxide dropwise until in excess.

Observations	Inferences
. Freekcsep	
Jisit water	
x Papers	
(1 mark)	(1 mark)

(ii) To the second portion in test tube, add aqueous ammonia solution dropwise until in excess.

	Observations	Inferences
More		
\$ <sup>0<sup>°</sup></sup>		
	(1 mark)	(1 mark)

(iii) To the third portion in a test-tube, add lead (II) nitrate solution and then warm the mixture.

Observations	Inferences
(1	
(1 mark)	(1 mark)